

COMULTIPLICATIVITY OF THE OZSVÁTH-SZABÓ CONTACT INVARIANT

JOHN A. BALDWIN

ABSTRACT. Suppose that S is a surface with boundary and that g and h are diffeomorphisms of S which restrict to the identity on the boundary. Let Y_g , Y_h , and Y_{hg} be the three-manifolds with open book decompositions given by (S, g) , (S, h) , and (S, hg) , respectively. We show that the Ozsváth-Szabó contact invariant is natural under a comultiplication map $\tilde{\mu} : \widehat{HF}(-Y_{hg}) \rightarrow \widehat{HF}(-Y_g) \otimes \widehat{HF}(-Y_h)$. It follows that if the contact invariants associated to the open books (S, g) and (S, h) are non-zero then the contact invariant associated to the open book (S, hg) is also non-zero. We extend this comultiplication to a map on $HF^+(-Y_{hg})$, and as a result we obtain obstructions to the three-manifold Y_{hg} being an L -space. We also use this to find restrictions on contact structures which are compatible with planar open books.

1. INTRODUCTION

In 2002, Ozsváth and Szabó discovered an invariant of contact 3-manifolds constructed as follows [11]. Given a contact 3-manifold (Y, ξ) , we can find a compatible (in the sense of Giroux [5]) open book decomposition (S, g) of Y with connected binding $K \subset Y$. If the genus of S is k then the knot Floer homology, $\widehat{HFK}(-Y, K, -k) \cong \mathbb{Z}$. Moreover, it is possible to find a Heegaard diagram for which the knot Floer homology in this $-k$ th filtration level is generated by an element of the chain complex $\widehat{CFK}(-Y, K)$ which also represents a cycle in the chain complex $\widehat{CF}(-Y)$. The contact invariant $c(\xi)$ is defined to be the image of this cycle in $\widehat{HF}(-Y)$.

The class $c(\xi)$ is well-defined up to sign (we use $\mathbb{Z}_2$ coefficients throughout to avoid ambiguity in sign), and it is an invariant of ξ up to isotopy of the contact structure. This invariant encodes information related to the tightness of ξ . For instance, Ozsváth and Szabó prove that if ξ is overtwisted, then $c(\xi) = 0$. On the other hand, if ξ is Stein fillable or strongly fillable, then $c(\xi) \neq 0$ [11], [10]. In a previous paper we show, in the case of contact structures compatible with genus one, one boundary component open books, that $c(\xi) = 0$ if and only if ξ is overtwisted for all but a small family of open books with reducible monodromies [1]. Honda, Kazez, and Matić have since shown this to be the case for all such open books [7], [6], [8]. However, the precise relationship between $c(\xi)$ and the tightness of ξ is still unknown – there are tight contact structures with vanishing contact invariant [4]. In fact, Lisca and Stipsicz conjecture that the contact invariant vanishes for contact structures with positive *Giroux torsion* [9].

As the contact invariant is defined in terms of a compatible open book decomposition, we often denote $c(\xi)$ by $c(S, g)$. This class satisfies the following naturality property [11]:

Theorem 1.1 (Ozsváth-Szabó). *If (S, g) is an open book decomposition for Y , and $\gamma \subset Y - L$ is a curve supported in a page of the open book (L is the binding), which is not homotopic to the boundary, then $(S, t_\gamma^{-1}g)$ induces an open book decomposition of $Y_{+1}(\gamma)$ (here, t_γ denotes the right-handed Dehn twist around γ). And under the map*

$$F_W : \widehat{HF}(-Y) \longrightarrow \widehat{HF}(-Y_{+1}(\gamma))$$

obtained by the two-handle addition (and summing over all spin^c structures), we have that

$$F_W(c(S, g)) = \pm c(S, t_\gamma^{-1}g).$$

In general, it is an open question as to whether tightness of the contact structures compatible with open books (S, g) and (S, h) implies tightness of the contact structure compatible with (S, hg) . This question is open even when h is a right-handed Dehn twist t_γ around a curve $\gamma \subset S$ [3]. Along these lines, however, Theorem 1.1 implies that if $c(S, g) \neq 0$, then $c(S, t_\gamma \circ g) \neq 0$. This behavior is generalized in Theorem 1.2, which follows from our main result.

Theorem 1.2. *Suppose that g and h are diffeomorphisms of S which restrict to the identity on ∂S and that $c(S, g) \neq 0$ and $c(S, h) \neq 0$. Then $c(S, hg) \neq 0$.*

This theorem has the immediate corollary:

Corollary 1.3. *If the contact structures compatible with (S, g) and (S, h) are strongly fillable, then the contact structure compatible with (S, hg) is tight.*

Another formulation of Theorem 1.2 is the statement that for a fixed S the set

$$N_S = \{g \in \text{Aut}(S, \partial S) \mid c(S, g) \neq 0\}$$

is a monoid under composition of diffeomorphisms, where here $\text{Aut}(S, \partial S)$ denotes the set of isotopy classes of diffeomorphisms of S that restrict to the identity on ∂S .

The key to this theorem is the observation that the contact invariant satisfies the naturality property below. For any g and h we exhibit a cobordism $X_{g,h}$ with

$$\partial X_{g,h} = -Y_g - Y_h + Y_{hg}$$

where Y_ϕ is the three-manifold with open book decomposition (S, ϕ) . This cobordism induces a chain map (multiplication)

$$m : \widehat{CF}(Y_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h) \rightarrow \widehat{CF}(Y_{hg}).$$

If we apply the $\text{Hom}_{\mathbb{Z}_2}(-, \mathbb{Z}_2)$ functor to this expression, we obtain a chain map (comultiplication)

$$\mu : \widehat{CF}(-Y_{hg}) \rightarrow \widehat{CF}(-Y_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(-Y_h).$$

We show that the contact invariants are natural under the corresponding map

$$\tilde{\mu} : \widehat{HF}(-Y_{hg}) \rightarrow \widehat{HF}(-Y_g) \otimes_{\mathbb{Z}_2} \widehat{HF}(-Y_h)$$

induced on homology. That is

Theorem 1.4. *The map $\tilde{\mu}$ takes $c(S, hg) \mapsto c(S, g) \otimes c(S, h)$.*

Hence, if $c(S, hg) = 0$ then either $c(S, g) = 0$ or $c(S, h) = 0$, and Theorem 1.2 follows immediately.

In Section 4, we generalize the result of Theorem 1.4 by examining analogous maps on HF^+ . We use this generalization to prove the following theorem.

Theorem 1.5. *If $c^+(S, hg) \in U^d \cdot HF^+(-Y_{hg})$, then $c^+(S \#_b S, g \# h) \in U^d \cdot HF^+(-(Y_g \# Y_h))$.*

In Theorem 1.5, $c^+(S, \phi)$ denotes the image in $HF^+(-Y_\phi)$ of $c(S, \phi)$ under the natural map $\widehat{HF}(-Y_\phi) \rightarrow HF^+(-Y_\phi)$. Furthermore, if (S, g) and (S, h) are open books compatible with contact structures (Y_g, ξ_g) and (Y_h, ξ_h) , respectively, then $(S \#_b S, g \# h)$ is an open book compatible with the contact structure $(Y_g \# Y_h, \xi_g \# \xi_h)$, where $\#_b$ denotes boundary connected sum.

The following corollaries of Theorem 1.5 provide obstructions to a contact three-manifold with open book (S, hg) having a compatible open book with planar pages. At the same time, we obtain obstructions to the three-manifold Y_{hg} being an L -space. The reader should compare these corollaries with those found by Ozsváth, Szabó, and Stipsicz in [13]. In what follows, $\mathfrak{s}(S, \phi)$ denotes the $spin^c$ structure associated to the contact 2-plane field which is compatible with (S, ϕ) .

Corollary 1.6. *Suppose that $c^+(S \#_b S, g \# h) \neq 0$ and that $c_1(\mathfrak{s}(S, g))$ is non-torsion. Then Y_{hg} is not an L -space and the contact structure corresponding to (S, hg) is not compatible with a planar open book.*

Corollary 1.7. *Suppose that (S, g) and (S, h) correspond to Stein fillable contact manifolds (Y_g, ξ_g) and (Y_h, ξ_h) with fillings (X_g, J_g) and (X_h, J_h) . Suppose further that $c_1(\mathfrak{s}(\xi_g)) = 0 = c_1(\mathfrak{s}(\xi_h))$ and $c_1(X_g, J_g) \neq 0$. Then Y_{hg} is not an L -space and the contact structure corresponding to (S, hg) is not compatible with a planar open book.*

1.1. Acknowledgements. I would like to thank Ko Honda for bringing to my attention the possibility of Theorem 1.2. I also wish to thank John Etnyre, Danny Gillam, Robert Lipshitz, and Shaffiq Welji for very helpful discussions. And, as always, I am indebted to Peter Ozsváth for his invaluable comments and suggestions.

2. HEEGAARD DIAGRAMS AND THE CONTACT CLASS

Honda, Kazez, and Matić give another interpretation of the Ozsváth-Szabó contact class in [7]. We use their reformulation in our proof of Theorem 1.2. Recall that the open book (S, g) is a decomposition of the 3-manifold $Y = S \times [0, 1] / \sim$, where $\sim$ is the relation defined by

$$\begin{aligned} (x, 1) &\sim (g(x), 0), & x \in S \\ (x, t) &\sim (x, s), & x \in \partial S, t, s \in [0, 1] \end{aligned}$$

Y has a Heegaard splitting $Y = H_1 \cup H_2$, where H_1 is the handlebody $S \times [0, 1/2]$ and H_2 is the handlebody $S \times [1/2, 1]$. Let S_t denote the page $S \times \{t\}$. If S has n boundary components and genus k then the Heegaard surface in this splitting is the genus $2k + n - 1$ surface $\Sigma = S_{1/2} \cup -S_0$. To specify a pointed Heegaard diagram for Y it remains to describe the α and β curves on Σ and the placement of a basepoint z . Choose $2k + n - 1$ disjoint

properly embedded arcs $a_1, \dots, a_{2k+n-1}$ on Σ so that $\Sigma \setminus \cup a_i$ is topologically a disk. For each i we obtain b_i by changing the arcs a_i via a small isotopy which moves the endpoints of the a_i along ∂S in the direction given by the orientation of ∂S so that a_i intersects b_i transversely in one point and with positive sign (where b_i inherits its orientation from a_i). See Figure 1 for an illustration of the a_i and b_i arcs on a surface S .




FIGURE 1. A surface S with multiple boundary components and genus greater than one. The a_i arcs are in red and the b_i arcs are in blue.

Now define

$$\begin{aligned}\alpha_i &= a_i \times \{1/2\} \cup a_i \times \{0\} \\ \beta_i &= b_i \times \{1/2\} \cup g(b_i) \times \{0\}.\end{aligned}$$

Place the basepoint z in the big region on $S_{1/2}$ (that is, not in one of the thin strip regions). For each $i = 1, \dots, 2k + n - 1$ let x_i be the intersection point on $S_{1/2}$ between α_i and β_i . Then $\overline{x_g} = \{x_1, \dots, x_{2k+n-1}\}$ is an intersection point between $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$ in $Sym^{2k+n-1}(\Sigma)$. Moreover, $\overline{x_g}$ is a cycle in $Hom(\widehat{CF}(Y), \mathbb{Z}_2) = \widehat{CF}(-Y)$ because of the placement of z . See Figure 2 for an illustration of the pointed Heegaard diagram for an open book.

Theorem 2.1 (Honda-Kazez-Matić). $[\overline{x_g}] \in \widehat{HF}(-Y)$ is the Ozsváth-Szabó contact class $c(S, g)$.

3. NATURALITY UNDER COMULTIPLICATION

Given a surface S with genus k and n boundary components, let a_i and b_i be the set of properly embedded arcs described above. We construct another set of disjoint properly embedded arcs c_i from the b_i by changing the arcs b_i via a small isotopy which moves the endpoints of the b_i along ∂S in the direction given by the orientation of ∂S . We require that both a_i and b_i intersect c_i transversely in one point and with positive sign (where c_i inherits its orientation from b_i). For any two diffeomorphisms g and h , we construct three sets of attaching curves on the Heegaard surface $\Sigma = S_{1/2} \cup -S_0$:

$$\begin{aligned}\alpha_i &= a_i \times \{1/2\} \cup a_i \times \{0\} \\ \beta_i &= b_i \times \{1/2\} \cup g(b_i) \times \{0\}\end{aligned}$$




FIGURE 2. The bottom figure is a pointed Heegaard diagram for the open book (S, g) , where S is a genus k surface with n boundary components and g is the composition of a left-handed Dehn twist around the curve p with a right-handed Dehn twist around the curve q . The α_i are in red and the β_i are in blue. We have also labeled the intersection point $\overline{x}_g = \{x_1, \dots, x_{2k+n-1}\}$.

$$\gamma_i = c_i \times \{1/2\} \cup h(g(c_i)) \times \{0\}.$$

Once again, we place the basepoint z in the big region of $S_{1/2}$ (outside of the thin strip regions). Then $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ is a pointed Heegaard triple-diagram and can be used as in [10] to construct a cobordism $X_{\alpha, \beta, \gamma}$ with

$$\partial X_{\alpha, \beta, \gamma} = -Y_{\alpha, \beta} - Y_{\beta, \gamma} + Y_{\alpha, \gamma}$$

where $Y_{\alpha, \beta}$ is the three manifold with Heegaard decomposition $(\Sigma, \bar{\alpha}, \bar{\beta})$ (and similarly for $Y_{\beta, \gamma}$ and $Y_{\alpha, \gamma}$). Such a cobordism induces a chain map

$$\widehat{CF}(Y_{\alpha, \beta}) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_{\beta, \gamma}) \rightarrow \widehat{CF}(Y_{\alpha, \gamma}).$$

By the description of the Heegaard diagram associated to an open book in section 2, it is clear that

$$Y_{\alpha,\beta} = Y_g$$

$$Y_{\beta,\gamma} = Y_h$$

$$Y_{\alpha,\gamma} = Y_{hg}.$$

Thus, we have a chain map

$$(1) \quad m : \widehat{CF}(Y_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h) \rightarrow \widehat{CF}(Y_{hg}).$$

If the pointed Heegaard triple-diagram $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ is *weakly-admissible* then this map is defined on the generators of $\widehat{CF}(Y_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h)$ by

$$m(\bar{a} \otimes \bar{b}) = \sum_{\bar{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\gamma} \sum_{\{\phi \in \pi_2(\bar{a}, \bar{b}, \bar{x}) \mid \mu(\phi)=0, n_z(\phi)=0\}} (\#\mathcal{M}(\phi))\bar{x}.$$

In this sum, $\pi_2(\bar{a}, \bar{b}, \bar{x})$ is the set of homotopy classes of *Whitney triangles* connecting $\bar{a}$, $\bar{b}$, and $\bar{x}$; $\mu(\phi)$ is the expected dimension of holomorphic representatives of ϕ ; $n_z(\phi)$ is the algebraic intersection number of ϕ with the subvariety $\{z\} \times \text{Sym}^{2k+n-2}(\Sigma) \subset \text{Sym}^{2k+n-1}(\Sigma)$; and $\mathcal{M}(\phi)$ is the moduli space of holomorphic representatives of ϕ . For more details, see [10].

As alluded to in the introduction, we apply the $\text{Hom}_{\mathbb{Z}_2}(-, \mathbb{Z}_2)$ functor to the expression in equation 1. If we represent each chain complex diagrammatically by drawing an arrow from x to y whenever y is a term in ∂x or when y is a term in the image of x under the map m , then applying the $\text{Hom}_{\mathbb{Z}_2}(-, \mathbb{Z}_2)$ functor corresponds to reversing the direction of every arrow. Doing so, we obtain a chain map

$$\mu : \widehat{CF}(-Y_{hg}) \rightarrow \widehat{CF}(-Y_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(-Y_h).$$

An element in $\widehat{CF}(-Y_{hg})$ is a sum of intersection points $\bar{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\gamma$. On such an intersection point the chain map μ is defined by

$$\mu(\bar{x}) = \sum_{\bar{a} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta, \bar{b} \in \mathbb{T}_\beta \cap \mathbb{T}_\gamma} \sum_{\{\phi \in \pi_2(\bar{a}, \bar{b}, \bar{x}) \mid \mu(\phi)=0, n_z(\phi)=0\}} (\#\mathcal{M}(\phi))(\bar{a} \otimes \bar{b})$$

as long as the pointed Heegaard triple-diagram $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ is weakly-admissible. To prove Theorem 1.4, we show that

$$\mu(\overline{x_{hg}}) = \overline{x_g} \otimes \overline{x_h}.$$

We complete the proof in two steps. First we show that $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ is weakly-admissible. Then we show that there is only one pair $\{\bar{a} \in \mathbb{T}_{\alpha,\beta}, \bar{b} \in \mathbb{T}_{\beta,\gamma}\}$ for which there exists a homotopy class $\phi \in \pi_2(\bar{a}, \bar{b}, \overline{x_{hg}})$ with $n_z(\phi) = 0$ and such that ϕ has a holomorphic representative. Moreover, $\bar{a} = \overline{x_g}$, $\bar{b} = \overline{x_h}$, and the number of holomorphic representatives of ϕ is one.

3.1. Weak Admissibility. We begin with two definitions from [10].

Definition 3.1. For a pointed Heegaard triple-diagram $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$, let $D_1, \dots, D_m$ be the connected regions of $\Sigma - \cup \alpha_i - \cup \beta_i - \cup \gamma_i$. And let $\psi = \sum_i p_i D_i$ be a formal linear combination of the D_i so that $n_z(\psi) = 0$ and $\partial\psi = \sum_i p_i \partial D_i$ is a linear combination of complete α , β , and γ curves. Then ψ is called a triply-periodic domain.

Definition 3.2. The pointed Heegaard triple-diagram is said to be weakly-admissible if every non-trivial triply-periodic domain ψ has both positive and negative coefficients.

For each $i = 1, \dots, 2k + n - 1$ the curves α_i , β_i , and γ_i intersect on $S_{1/2}$ in the arrangement depicted in Figure 3.




FIGURE 3. The curve α_i is in red, β_i in blue, and γ_i in yellow. We have labeled regions $D_1, \dots, D_6$ as well as the i th components of the various contact classes $\overline{x_g}$, $\overline{x_h}$, and $\overline{x_{hg}}$.

If $\psi = \sum_j p_j D_j$ is a triply-periodic domain, then $p_6 = 0$ since D_6 contains the basepoint z . Since $\partial\psi$ includes some number of complete α_i curves,

$$p_2 = p_3 - p_4 = -p_5.$$

Therefore, ψ has both positive and negative coefficients unless $p_2 = p_5 = 0$ and $p_3 = p_4$. So let's assume the latter. Since $\partial\psi$ includes some number of complete β_i curves,

$$p_1 = -p_3 = 0.$$

So, either ψ has both positive and negative coefficients or $p_1 = \dots = p_6 = 0$ and $\partial\psi$ includes no α_i , β_i , or γ_i curves. If we carry out this analysis for all $i = 1, \dots, 2k + n - 1$ we see that either ψ has both positive and negative coefficients or else it is trivial. Hence $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ is weakly-admissible. $\square$

3.2. Completing the proof of Theorem 1.4. Let Δ denote the 2-simplex and label its vertices clockwise $v_\alpha, v_\beta, v_\gamma$. Let e_α be the edge opposite v_α (and similarly for e_β and e_γ). The boundary of Δ inherits the standard counterclockwise orientation. Then

Definition 3.3. A map $u : \Delta \rightarrow \text{Sym}^{2k+n-1}(\Sigma)$ satisfying $u(v_\gamma) = \bar{a}$, $u(v_\alpha) = \bar{b}$, and $u(v_\beta) = \overline{x_{hg}}$, and $u(e_\alpha) \subset \mathbb{T}_\alpha$, $u(e_\beta) \subset \mathbb{T}_\beta$, and $u(e_\gamma) \subset \mathbb{T}_\gamma$ is called a Whitney triangle between $\bar{a}$, $\bar{b}$, and $\overline{x_{hg}}$. This map u is represented schematically in Figure 4.




FIGURE 4.

We can represent $\phi \in \pi_2(\bar{a}, \bar{b}, \overline{x_{hg}})$ by a 2-chain $\hat{\phi} = \sum_j p_j D_j$ whose oriented boundary consists of α arcs from $\bar{a}$ to $\overline{x_{hg}}$, β arcs from $\bar{b}$ to $\bar{a}$, and γ arcs from $\overline{x_{hg}}$ to $\bar{b}$. Suppose $n_z(\phi) = 0$ and ϕ has a holomorphic representative. Then $n_z(\hat{\phi}) = 0$ and the p_j are all non-negative. We refer to Figure 3 for the local picture near the i th component of the contact classes $\overline{x_g}$, $\overline{x_h}$, and $\overline{x_{hg}}$. Write

$$\hat{\phi} = p_1 D_1 + \cdots + p_6 D_6 + \sum_{j>6} p_j D_j.$$

Now we can analyze the possibilities for $p_1, \dots, p_6$ given the boundary constraints on $\hat{\phi}$. $(\overline{x_{hg}})_i$ must be a corner of the region defined by $\hat{\phi}$; moreover this corner is such that we enter $(\overline{x_{hg}})_i$ along an arc of α_i and we leave along an arc of γ_i . Therefore, $p_6 + p_3 = p_2 + p_4 + 1$. If $(\overline{x_h})_i$ is not a corner, then $p_3 + p_1 = p_2 + p_6$. Note that $p_6 = 0$ since $n_z(\hat{\phi}) = 0$. Thus, these two equations become

$$\begin{aligned} p_3 &= p_2 + p_4 + 1 \\ p_3 + p_1 &= p_2. \end{aligned}$$

Subtracting the second equation from the first, we have

$$-p_1 = p_4 + 1$$

which implies that either p_1 or p_4 is negative, which cannot happen since ϕ has a holomorphic representative. Therefore, $(\overline{x_h})_i$ is a corner. The same type of analysis shows that $(\overline{x_g})_i$ is a corner.

Since $(\overline{x_h})_i$ is a corner, either $p_1 + p_3 + 1 = p_2$ or $p_1 + p_3 = p_2 + 1$. Substituting $p_3 = p_2 + p_4 + 1$ into both expressions, we have the two possibilities $p_1 + p_2 + p_4 + 2 = p_2$ or $p_1 + p_2 + p_4 = p_2$.

We can rule out the first possibility as it implies that either p_1 or p_4 is negative. And the second possibility holds only if $p_1 = p_4 = 0$. So, to summarize what we know so far: $p_1 = 0$, $p_3 = p_2 + 1$, $p_4 = 0$, and $p_6 = 0$.

Since $(\overline{x_g})_i$ is a corner, then either $p_5 + p_3 = p_4 + p_6 + 1$ or $p_5 + p_3 + 1 = p_4 + p_6$. Substituting what we know of p_3 , p_4 , and p_6 into these two expressions, we obtain the two possibilities $p_5 + p_2 + 1 = 1$ or $p_5 + p_2 + 2 = 0$. We can rule out the second possibility as it implies that either p_5 or p_2 is negative. And the first possibility holds only if $p_5 = p_2 = 0$. Thus, we have determined that the only possibility for the values $p_1, \dots, p_6$ are:

$$p_1 = p_2 = p_4 = p_5 = p_6 = 0$$

$$p_3 = 1.$$

Because the same analysis works for every $i = 1, \dots, 2k+n-1$ and because every component of $\partial\widehat{\phi}$ must contain some $(\overline{x_{hg}})_i$ we can conclude that $\widehat{\phi}$ is the linear combination which is the sum of precisely one of these small triangular regions (D_3 in figure 3) for each i . Therefore, any holomorphic triangle ϕ between $\bar{a}$, $\bar{b}$, and $\overline{x_{hg}}$ with $n_z(\phi) = 0$ is, in fact, a triangle between $\overline{x_g}$, $\overline{x_h}$, and $\overline{x_{hg}}$, and can be expressed as a product of these small triangles in our Heegaard diagram. Moreover, since each of these disjoint triangular regions is topologically a disk, and we have specified the image of three boundary points, $\#\mathcal{M}(\phi) = 1$ by the Riemann Mapping Theorem. Hence,

$$\mu(\overline{x_{hg}}) = \overline{x_g} \otimes \overline{x_h}.$$

Therefore,

$$\tilde{\mu}(c(S, hg)) = c(S, g) \otimes c(S, h)$$

and the proof of Theorem 1.4 is complete. As mentioned in the Introduction, Theorem 1.2 follows immediately. $\square$

4. A GENERALIZATION OF THEOREM 1.4

4.1. $CF^{\leq 0}$ and connected sums. For a $spin^c$ structure $\mathfrak{s}$ on Y and a pointed Heegaard diagram $(\Sigma, \bar{\alpha}, \bar{\beta}, z)$ for Y which is *strongly $\mathfrak{s}$ -admissible*, recall that we can define a chain complex $CF^{\leq 0}(Y, \mathfrak{s})$ which is finitely generated as a $\mathbb{Z}_2[U]$ module [10]. The generators of $CF^{\leq 0}(Y, \mathfrak{s})$ are pairs of the form $[\bar{x}, i]$, where $\bar{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$, $\mathfrak{s}_z(\bar{x}) = \mathfrak{s}$, $i \in \mathbb{Z}^{\leq 0}$, and U acts by $U \cdot [\bar{x}, i] = [\bar{x}, i - 1]$. The differential on $CF^{\leq 0}(Y, \mathfrak{s})$ is given by

$$\partial[\bar{x}, i] = \sum_{\bar{y} \in \mathbb{T}_\alpha \cap \mathbb{T}_\gamma, \{\phi \in \pi_2(\bar{x}, \bar{y}) \mid \mu(\phi)=1\}} \sum \#(\frac{\mathcal{M}(\phi)}{\mathbb{R}}) \cdot [\bar{y}, i - n_z(\phi)].$$

We can identify $\widehat{CF}(Y, \mathfrak{s})$ with $CF^{\leq 0}(Y, \mathfrak{s})/U \cdot CF^{\leq 0}(Y, \mathfrak{s})$, so there is a natural quotient map

$$\pi : CF^{\leq 0}(Y, \mathfrak{s}) \rightarrow \widehat{CF}(Y, \mathfrak{s}).$$

If A is a $\mathbb{Z}_2[U]$ module, let $A^\vee$ denote $\text{Hom}_{\mathbb{Z}_2[U]}(A, \mathbb{Z}_2[U, U^{-1}]/\mathbb{Z}_2[U])$. Then observe that we can identify $CF^{\leq 0}(Y, \mathfrak{s})^\vee$ with $CF^+(-Y, \mathfrak{s})$, and $\widehat{CF}(Y, \mathfrak{s})^\vee$ with $\widehat{CF}(-Y, \mathfrak{s})$ as complexes over

$\mathbb{Z}_2[U]$ and $\mathbb{Z}_2$, respectively.¹ In fact, applying the $\text{Hom}_{\mathbb{Z}_2[U]}(-, \mathbb{Z}_2[U, U^{-1}]/\mathbb{Z}_2[U])$ functor to the expression above, we obtain the natural inclusion map

$$\pi^\vee : \widehat{CF}(-Y, \mathfrak{s}) \rightarrow CF^+(-Y, \mathfrak{s})$$

which sends

$$\bar{x} \mapsto [\bar{x}, 0]$$

for $\bar{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$.

In [12] the authors construct a homotopy equivalence

$$f^{\leq 0} : CF^{\leq 0}(Y_1, \mathfrak{s}_1) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_2, \mathfrak{s}_2) \rightarrow CF^{\leq 0}(Y_1 \# Y_2, \mathfrak{s}_1 \# \mathfrak{s}_2)$$

as follows. Let $(\Sigma_i, \bar{\alpha}_i, \bar{\beta}_i, z_i)$ be a strongly $\mathfrak{s}_i$ -admissible pointed Heegaard diagram for Y_i , for $i = 1, 2$. Consider the pointed Heegaard triple-diagram $(\Sigma_1 \# \Sigma_2, \bar{\alpha}_1 \cup \bar{\alpha}'_2, \bar{\beta}_1 \cup \bar{\alpha}_2, \bar{\beta}'_1 \cup \bar{\alpha}_2, z)$, where the connected sum of Σ_1 with Σ_2 occurs at the z_i , z is a point in the connected sum region, and $\bar{\alpha}'_2$ and $\bar{\beta}'_1$ are exact Hamiltonian translates of $\bar{\alpha}_2$ and $\bar{\beta}_1$ so that this new diagram is admissible. Suppose that the genera of Σ_1 and Σ_2 are g_1 and g_2 and let $\Theta_1 \in \mathbb{T}_{\beta_1} \cap \mathbb{T}_{\beta'_1}$ and $\Theta_2 \in \mathbb{T}_{\alpha_2} \cap \mathbb{T}_{\alpha'_2}$ be the top graded intersection points in $\widehat{CF}(\#^{g_1}(S^1 \times S^2), \mathfrak{s}_0)$ and $\widehat{CF}(\#^{g_2}(S^1 \times S^2), \mathfrak{s}_0)$. Then the maps $[\bar{x}, i] \mapsto [\bar{x} \times \Theta_2, i]$ and $[\bar{y}, j] \mapsto [\Theta_1 \times \bar{y}, j]$ define chain maps

$$\Phi_1 : CF^{\leq 0}(Y_1, \mathfrak{s}_1) \rightarrow CF^{\leq 0}(Y_1 \# \#^{g_2}(S^1 \times S^2), \mathfrak{s}_1 \# \mathfrak{s}_0),$$

and

$$\Phi_2 : CF^{\leq 0}(Y_2, \mathfrak{s}_2) \rightarrow CF^{\leq 0}(\#^{g_2}(S^1 \times S^2) \# Y_2, \mathfrak{s}_0 \# \mathfrak{s}_2).$$

The pointed Heegaard triple-diagram above can be used to define a map

$$\begin{aligned} \Gamma : CF^{\leq 0}(Y_1 \# \#^{g_2}(S^1 \times S^2), \mathfrak{s}_1 \# \mathfrak{s}_0) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(\#^{g_2}(S^1 \times S^2) \# Y_2, \mathfrak{s}_0 \# \mathfrak{s}_2) \\ \longrightarrow CF^{\leq 0}(Y_1 \# Y_2, \mathfrak{s}_1 \# \mathfrak{s}_2). \end{aligned}$$

The map $f^{\leq 0}$ is then defined to be

$$f^{\leq 0} = \Gamma \circ (\Phi_1 \otimes \Phi_2).$$

The maps in this composition are all U -equivariant by construction; hence, so is $f^{\leq 0}$. In [12], Ozsváth and Szabó show that $f^{\leq 0}$ is a homotopy equivalence. We define a homotopy equivalence

$$f : \widehat{CF}(Y_1, \mathfrak{s}_1) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_2, \mathfrak{s}_2) \rightarrow \widehat{CF}(Y_1 \# Y_2, \mathfrak{s}_1 \# \mathfrak{s}_2)$$

in exactly the same way.²

¹We have already been identifying $\widehat{CF}(-Y)$ with $\text{Hom}_{\mathbb{Z}_2}(\widehat{CF}(Y), \mathbb{Z}_2)$, but the latter is isomorphic as a $\mathbb{Z}_2$ module to $\widehat{CF}(Y)^\vee$ since we are thinking of $\widehat{CF}(Y)$ as a $\mathbb{Z}_2[U]$ module where the action of U is multiplication by 0.

²In [12], the authors define a homotopy equivalence between these two chain complexes in a slightly different and more direct way. However, the map f defined here is better suited for our purposes.

4.2. HF^+ and the contact invariant. Let ϕ denote the Whitney triangle between $\overline{x_g}$, $\overline{x_h}$, and $\overline{x_{hg}}$ found in Section 3, and let $\mathfrak{s}(\phi)$ denote the $spin^c$ structure on $X_{g,h}$ corresponding to ϕ . Moreover, let $\mathfrak{s}_g$, $\mathfrak{s}_h$, and $\mathfrak{s}_{hg}$ denote the induced $spin^c$ structures on Y_g , Y_h , and Y_{hg} . Then there is a chain map

$$m_{\mathfrak{s}(\phi)} : \widehat{CF}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h, \mathfrak{s}_h) \rightarrow \widehat{CF}(Y_{hg}, \mathfrak{s}_{hg})$$

which is a refinement of the map m defined in Section 3. The difference between the two is that $m_{\mathfrak{s}(\phi)}$ counts only those Whitney triangles which correspond to the $spin^c$ structure $\mathfrak{s}(\phi)$. Applying the $\text{Hom}_{\mathbb{Z}_2[U]}(-, \mathbb{Z}_2[U, U^{-1}]/\mathbb{Z}_2[U])$ functor as before and taking homology, it is clear that the induced map

$$(m_{\mathfrak{s}(\phi)}^\vee)_* : \widehat{HF}(-Y_{hg}, \mathfrak{s}_{hg}) \rightarrow \widehat{HF}(-Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2} \widehat{HF}(-Y_h, \mathfrak{s}_h)$$

still takes $c(S, hg) \mapsto c(S, g) \otimes c(S, h)$. Mirroring the notation in Section 3, we denote this map by $\tilde{\mu}_{\mathfrak{s}(\phi)}$.

The pointed Heegaard triple-diagram $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ from Section 3 also gives a U -equivariant chain map

$$m_{\mathfrak{s}(\phi)}^{\leq 0} : CF^{\leq 0}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_h, \mathfrak{s}_h) \rightarrow CF^{\leq 0}(Y_{hg}, \mathfrak{s}_{hg}).^3$$

Let

$$\pi_{g \otimes h} : CF^{\leq 0}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_h, \mathfrak{s}_h) \rightarrow \widehat{CF}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h, \mathfrak{s}_h)$$

and

$$\pi_{hg} : CF^{\leq 0}(Y_{hg}, \mathfrak{s}_{hg}) \rightarrow \widehat{CF}(Y_{hg}, \mathfrak{s}_{hg})$$

denote the quotient maps discussed in Subsection 4.1. Then the following diagram commutes.

$$\begin{array}{ccc} CF^{\leq 0}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_h, \mathfrak{s}_h) & \xrightarrow{m_{\mathfrak{s}(\phi)}^{\leq 0}} & CF^{\leq 0}(Y_{hg}, \mathfrak{s}_{hg}) \\ \downarrow \pi_{g \otimes h} & & \downarrow \pi_{hg} \\ \widehat{CF}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h, \mathfrak{s}_h) & \xrightarrow{m_{\mathfrak{s}(\phi)}} & \widehat{CF}(Y_{hg}, \mathfrak{s}_{hg}) \end{array}$$

After applying the $\text{Hom}_{\mathbb{Z}_2[U]}(-, \mathbb{Z}_2[U, U^{-1}]/\mathbb{Z}_2[U])$ functor and taking homology, we obtain the commutative diagram

³For this map to be well-defined, we need $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ to be strongly $\mathfrak{s}(\phi)$ -admissible. We can assume, however, that this is the case since the Heegaard diagram $(\Sigma, \bar{\alpha}, \bar{\beta}, \bar{\gamma}, z)$ can be made strongly $\mathfrak{s}(\phi)$ -admissible by winding the $\bar{\alpha}$, $\bar{\beta}$, and $\bar{\gamma}$ circles around curves contained strictly in the $-S_0$ portion of the surface Σ [10].

$$\begin{array}{ccc}
H_*((CF^{\leq 0}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_h, \mathfrak{s}_h))^{\vee}) & \xleftarrow{\tilde{\mu}_{\mathfrak{s}(\phi)}^+} & HF^+(-Y_{hg}, \mathfrak{s}_{hg}) \\
\uparrow (\pi_{g \otimes h}^{\vee})_* & & \uparrow (\pi_{hg}^{\vee})_* \\
\widehat{HF}(-Y_g, \mathfrak{s}_g) \otimes \widehat{HF}(-Y_h, \mathfrak{s}_h) & \xleftarrow{\tilde{\mu}_{\mathfrak{s}(\phi)}} & \widehat{HF}(-Y_{hg}, \mathfrak{s}_{hg})
\end{array}$$

where $\tilde{\mu}_{\mathfrak{s}(\phi)}^+ = ((m_{\mathfrak{s}(\phi)}^{\leq 0})^{\vee})_*$. Recall that for a contact three-manifold (Y, ξ) , the class $c^+(\xi) \in HF^+(-Y)$ is defined to be the image of $c(\xi)$ under the natural map $\widehat{HF}(-Y) \rightarrow HF^+(-Y)$ [13]. As was mentioned in Subsection 4.1, $(\pi_{hg}^{\vee})_*$ is this natural map, and therefore

$$(\pi_{hg}^{\vee})_*(c(S, hg)) = c^+(S, hg).$$

Let $K_{g \otimes h}$ denote $\tilde{\mu}_{\mathfrak{s}(\phi)}^+(c^+(S, hg))$. Then by the commutativity of this diagram,

$$K_{g \otimes h} = \tilde{\mu}_{\mathfrak{s}(\phi)}^+(c^+(S, hg)) = (\pi_{g \otimes h}^{\vee})_*(c(S, g) \otimes c(S, h)).$$

Returning to our discussion of connected sums, let

$$\pi_{g \# h} : CF^{\leq 0}(Y_g \# Y_h, \mathfrak{s}_g \# \mathfrak{s}_h) \rightarrow \widehat{CF}(Y_g \# Y_h, \mathfrak{s}_g \# \mathfrak{s}_h)$$

denote the natural quotient map, and let $f^{\leq 0}$ and f be the homotopy equivalences described in Subsection 4.1. Then the diagram below commutes.

$$\begin{array}{ccc}
CF^{\leq 0}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_h, \mathfrak{s}_h) & \xrightarrow{f^{\leq 0}} & CF^{\leq 0}(Y_g \# Y_h, \mathfrak{s}_g \# \mathfrak{s}_h) \\
\downarrow \pi_{g \otimes h} & & \downarrow \pi_{g \# h} \\
\widehat{CF}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2} \widehat{CF}(Y_h, \mathfrak{s}_h) & \xrightarrow{f} & \widehat{CF}(Y_g \# Y_h, \mathfrak{s}_g \# \mathfrak{s}_h)
\end{array}$$

Again, after applying the $\text{Hom}_{\mathbb{Z}_2[U]}(-, \mathbb{Z}_2[U, U^{-1}]/\mathbb{Z}_2[U])$ functor and taking homology, we obtain the commutative diagram

$$\begin{array}{ccc}
H_*((CF^{\leq 0}(Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2[U]} CF^{\leq 0}(Y_h, \mathfrak{s}_h))^{\vee}) & \xleftarrow{\tilde{k}^+} & HF^+(-(Y_g \# Y_h), \mathfrak{s}_g \# \mathfrak{s}_h) \\
\uparrow (\pi_{g \otimes h}^{\vee})_* & & \uparrow (\pi_{g \# h}^{\vee})_* \\
\widehat{HF}(-Y_g, \mathfrak{s}_g) \otimes_{\mathbb{Z}_2} \widehat{HF}(-Y_h, \mathfrak{s}_h) & \xleftarrow{\tilde{k}} & \widehat{HF}(-(Y_g \# Y_h), \mathfrak{s}_g \# \mathfrak{s}_h)
\end{array}$$

where $\tilde{k}^+ = ((f^{\leq 0})^{\vee})_*$ and $\tilde{k} = (f^{\vee})_*$. At this point, we are ready to generalize Theorem 1.4.

Theorem 4.1. *The map $(\tilde{k}^+)^{-1} \circ \tilde{\mu}_{\mathfrak{s}(\phi)}^+ : HF^+(-Y_{hg}, \mathfrak{s}_{hg}) \rightarrow HF^+(-(Y_g \# Y_h), \mathfrak{s}_g \# \mathfrak{s}_h)$ is U -equivariant and takes $c^+(S, hg) \mapsto c^+(S \#_b S, g \# h)$.*

Observe that Theorem 1.5 follows immediately as a corollary. Recall from the introduction that the contact structure compatible with the open book $(S \#_b S, g \# h)$ is the connected sum of the contact structures compatible with (S, g) and (S, h) .

Proof of Theorem 4.1. The maps $\tilde{\mu}_{\mathfrak{s}(\phi)}^+$ and $(\tilde{k}^+)^{-1}$ are certainly U -equivariant, so their composition is as well. Moreover, $\tilde{k} = (f^\vee)_*$ takes $c(S \#_b S, g \# h) \mapsto c(S, g) \otimes c(S, h)$. This follows from precisely the same sort of argument as was used in Section 3 to show that $\tilde{\mu} = (m^\vee)_*$ takes $c(S, hg) \mapsto c(S, g) \otimes c(S, h)$. In the pointed Heegaard triple-diagram $(\Sigma_1 \# \Sigma_2, \bar{\alpha}_1 \cup \bar{\alpha}'_2, \bar{\beta}_1 \cup \bar{\alpha}_2, \bar{\beta}'_1 \cup \bar{\alpha}_2, z)$ used to construct the map Γ , the only holomorphic Whitney triangle $\phi \in \pi_2(\bar{a}, \bar{b}, \bar{x}_g \times \bar{x}_g)$ with $n_z(\phi) = 0$ is a product of small triangles connecting $\bar{x}_g \times \Theta_2$, $\Theta_1 \times \bar{x}_h$, and $\bar{x}_g \times \bar{x}_h$. Therefore, the map $f^\vee$ takes $\bar{x}_g \times \bar{x}_h \mapsto \bar{x}_g \otimes \bar{x}_h$ on the level of chains, and $(f^\vee)_*$ takes $c(S \#_b S, g \# h) \mapsto c(S, g) \otimes c(S, h)$.

Hence,

$$((\pi_{g \otimes h}^\vee)_* \circ \tilde{k})(c(S \#_b S, g \# h)) = (\pi_{g \otimes h}^\vee)_*(c(S, g) \otimes c(S, h)) = K_{g \otimes h}.$$

Therefore, by commutativity,

$$K_{g \otimes h} = (\tilde{k}^+ \circ (\pi_{g \# h}^\vee)_*)(c(S \#_b S, g \# h)) = \tilde{k}^+(c^+(S \#_b S, g \# h)).$$

Thus, $(\tilde{k}^+)^{-1}(K_{g \otimes h}) = c^+(S \#_b S, g \# h)$. But this implies that

$$((\tilde{k}^+)^{-1} \circ \tilde{\mu}_{\mathfrak{s}(\phi)}^+)(c^+(S, hg)) = (\tilde{k}^+)^{-1}(K_{g \otimes h}) = c^+(S \#_b S, g \# h).$$

□

5. L -SPACES AND PLANAR OPEN BOOKS

Etnyre recently showed that while every overtwisted contact structure has a compatible open book with planar pages, there are fillable contact structures which do not [2]. More recently, Ozsváth, Szabó, and Stipsicz found a Heegaard Floer homology obstruction to a contact structure having a compatible open book with planar pages, and were able to reproduce some of Etnyre's results [13]. Their main result is the following:

Theorem 5.1 (Ozsváth-Szabó-Stipsicz). *If the contact three-manifold (Y, ξ) has a compatible open book with planar pages then $c^+(\xi) \in U^d \cdot HF^+(-Y)$ for all $d \in \mathbb{N}$.*

They prove the following corollaries which are based upon this principle.

Corollary 5.2 (Ozsváth-Szabó-Stipsicz). *Suppose that $c^+(\xi) \neq 0$ and $c_1(\mathfrak{s}(\xi))$ is non-torsion. Then it cannot be the case that $c^+(\xi) \in U^d \cdot HF^+(-Y)$ for all $d \in \mathbb{N}$. In particular, (Y, ξ) is not supported by a planar open book.*

Corollary 5.3 (Ozsváth-Szabó-Stipsicz). *Suppose that the contact three-manifold (Y, ξ) has a Stein filling (X, J) with $c_1(\mathfrak{s}(\xi)) = 0$ and $c_1(X, J) \neq 0$. Then it cannot be the case that $c^+(\xi) \in U^d \cdot HF^+(-Y)$ for all $d \in \mathbb{N}$. In particular, (Y, ξ) is not supported by a planar open book.*

Our Corollaries 1.6 and 1.7 now follow from Theorems 1.5, 4.1, and the above corollaries of Ozsváth, Szabó, and Stipsicz.

Proof of Corollary 1.6. If $c_1(\mathfrak{s}(S, g))$ is non-torsion, then so is

$$c_1(\mathfrak{s}(S \#_b S, g \# h)) = c_1(\mathfrak{s}(S, g)) \oplus c_1(\mathfrak{s}(S, h)).$$

If, in addition, $c^+(S \#_b S, g \# h) \neq 0$, then Corollary 5.2 implies that it cannot be the case that $c^+(S \#_b S, g \# h) \in U^d \cdot HF^+(-(Y_g \# Y_h))$ for all $d \in \mathbb{N}$. Thus, Theorem 1.5 demands that it cannot be the case that $c^+(S, hg) \in U^d \cdot HF^+(-Y_{hg})$ for all $d \in \mathbb{N}$. It follows that Y_{hg} cannot be an L -space. Furthermore, it follows from Theorem 5.1 of Ozsváth, Szabó, and Stipsicz that the contact structure supported by (S, hg) is not compatible with a planar open book. $\square$

Proof of Corollary 1.7. If (S, g) and (S, h) correspond to contact manifolds (Y_g, ξ_g) and (Y_h, ξ_h) with Stein fillings (X_g, J_g) and (X_h, J_h) , then $(S \#_b S, g \# h)$ corresponds to the contact manifold $(Y_g \# Y_h, \xi_g \# \xi_h)$ with Stein filling $(X_g \#_b X_h, J_g \#_b J_h)$. If $c_1(\mathfrak{s}(\xi_g)) = 0$ and $c_1(\mathfrak{s}(\xi_h)) = 0$, then

$$c_1(\mathfrak{s}(\xi_g \# \xi_h)) = c_1(\mathfrak{s}(\xi_g)) \oplus c_1(\mathfrak{s}(\xi_h)) = 0.$$

Moreover, if $c_1(X_g, J_g) \neq 0$, then

$$c_1(X_g \#_b X_h, J_g \#_b J_h) = c_1(X_g, J_g) \oplus c_1(X_h, J_h) \neq 0.$$

Then, by Corollary 5.3, it cannot be the case that $c^+(S \#_b S, g \# h) \in U^d \cdot HF^+(-(Y_g \# Y_h))$ for all $d \in \mathbb{N}$. And, just as before, Theorem 1.5 then implies that it cannot be the case that $c^+(S, hg) \in U^d \cdot HF^+(-Y_{hg})$ for all $d \in \mathbb{N}$. It follows that Y_{hg} cannot be an L -space and that the contact structure supported by (S, hg) is not compatible with a planar open book. $\square$

It is not clear whether we can replace the condition that $c^+(S \#_b S, g \# h) \neq 0$ in the formulation of Corollary 1.6 by the condition that $c^+(S, g) \neq 0$ and $c^+(S, h) \neq 0$. For the contact invariant c defined in $\widehat{HF}(-Y)$, $c(S, g) \neq 0$ and $c(S, h) \neq 0$ implies that $c(S \#_b S, g \# h) \neq 0$. It is not immediately obvious that the same is true in general for the contact invariant c^+ . There are, however, *special* cases in which the same holds for c^+ . For instance, if (S, g) and (S, h) support strongly fillable contact structures, then so does $(S \#_b S, g \# h)$, and hence, $c^+(S \#_b S, g \# h) \neq 0$. More useful perhaps, is the following.

Claim 5.4. *If $c^+(S, g) \neq 0$ and (S, h) supports a Stein fillable contact structure, then $c^+(S \#_b S, g \# h) \neq 0$.*

Proof of Claim 5.4. If (S, h) supports a Stein fillable contact structure, then after a number of positive stabilizations (S, h) is equivalent to an open book (S', ϕ) , where ϕ is the composition of right-handed Dehn twists around curves in S' . Hence, $(S \#_b S, g \# h)$ is equivalent via the same number of positive stabilizations to the open book $(S \#_b S', g \# \phi)$. Therefore, by the naturality of the contact invariant c^+ under composition with left-handed Dehn twists, if $c^+(S \#_b S', g \# id) \neq 0$, then $c^+(S \#_b S', g \# \phi) \neq 0$.⁴ Let $m = 2k + r$, where k is the genus of

⁴Theorem 1.4 was stated only for the contact invariant c , but the same holds for c^+ via a commutative diagram chase.

S' and r is the number of boundary components of S' . Then $(S \#_b S', g \# id)$ is an open book for the manifold $Y_g \# \#^m(S^1 \times S^2)$. There is an isomorphism

$$HF^+(-(Y_g \# \#^m(S^1 \times S^2)), \mathfrak{s}_g \# \mathfrak{s}_0) \rightarrow HF^+(-Y_g, \mathfrak{s}_g) \wedge^* H^1(\#^m(S^1 \times S^2))$$

[12, Proposition 6.4]. It is clear from the construction that this isomorphism takes

$$c^+(S \#_b S', g \# id) \mapsto c^+(S, g) \wedge \Theta$$

where Θ is the lowest graded element of $\wedge^* H^1(\#^m(S^1 \times S^2))$. Therefore, $c^+(S, g) \neq 0$ if and only if $c^+(S \#_b S', g \# id) \neq 0$. Putting these facts together, we get the claim. $\square$

It remains to be seen what can be shown using these techniques. It would be very interesting, as Etnyre mentions in [2], to find a non-fillable contact structure which is not supported by an open book with planar pages. To this end it is enough to find, by Corollary 1.6, a Stein fillable open book (S, g) with $c_1(\mathfrak{s}(S, g))$ non-torsion and an open book (S, h) with $c^+(S, h) \neq 0$ such that (S, hg) is non-fillable.

REFERENCES

- [1] J. A. Baldwin. Tight contact structures and genus one fibered knots. 2006, math.SG/0604580.
- [2] J. B. Etnyre. Planar open book decompositions and contact structures. 2004, math.SG/0404267.
- [3] J. B. Etnyre. Lectures on open book decompositions and contact structures. 2005, math.SG/0409402.
- [4] P. Ghiggini. Ozsváth and Z. Szabó invariants and fillability of contact structures. 2005, math.GT/0403367.
- [5] E. Giroux. Convexité en topologie de contact. *Comment. Math. Helv.*, 66:637–677, 1991.
- [6] K. Honda, W. Kazez, and G. Matić. Right-veering diffeomorphisms of a compact surface with boundary I. 2005, math.GT/0510639.
- [7] K. Honda, W. Kazez, and G. Matić. On the contact class in Heegaard Floer homology. 2006, math.GT/0609734.
- [8] K. Honda, W. Kazez, and G. Matić. Right-veering diffeomorphisms of a compact surface with boundary II. 2006, math.GT/0603626.
- [9] P. Lisca and A. Stipsicz. Contact Ozsváth-Szabó invariants and Giroux torsion. 2006, math.SG/0604268.
- [10] P. Ozsváth and Z. Szabó. Holomorphic disks and topological invariants for closed three-manifolds. 2001, math.SG/0101206.
- [11] P. Ozsváth and Z. Szabó. Heegaard Floer homologies and contact structures. 2002, math.SG/0210127.
- [12] P. Ozsváth and Z. Szabó. Holomorphic disks and three-manifold invariants: properties and applications. 2003, math.SG/0105202.
- [13] P. Ozsváth, Z. Szabó, and A. I. Stipsicz. Planar open books and Floer homology. 2005, math.SG/0504403.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027

E-mail address: baldwin@math.columbia.edu